

Center for Women's Gastroenterology
Providence, RI
(401) 793-7080

Lifespan Physician's Group, Inc.

East Providence, RI (401)606-4260 Thomas DeNucci, MD Fadlallah Habr, MD Zilla Hussain, MD Sarah Hyder, MD Colleen Kelly, MD Amanda Pressman, MD Shannon Simmons, MD Laura Massa, NP Erica Taylor, PA Your procedure will take place at: ○ Rhode Island Hospital, 593 Eddy Street, Providence, RI - Main Building, 9th Floor Medical Office Center (MOC) Building, 2 Dudley Street, 3rd Floor, Providence, RI. Wayland Surgical Center, 17 Seekonk Street, Providence, RI The Miriam Hospital, 164 Summit Avenue, Providence, RI O Blackstone Surgical Center, 1526 Atwood Avenue, Suite #300, Johnston, RI

	On the ada			for Colonoscopy			
		· · · · ·	•		list of ALL your medic		
Due to unexpected emergencies, your procedure time cannot be guaranteed. Please contact your insurance company prior to your procedure to understand your responsibility for any co-payment and/or deductible. Note: This is a hospital-based Endoscopy Unit.							
Date:_	A	ppointment Time: _		Arrival Time:	With:	Dr	
Import	tant						
	you should take preparation ma A responsible a on the day of yo accompanied by Please do not w tant: Notify your p	the pill in the more y interfere with about dult must be availal our procedure. You y a responsible aduly ear perfumes or bout onysician if you are to	ning with your I sorption of the police to come into must have som It. ody lotions on the	ight breakfast and u pill. Resume your bi the Endoscopy suit eone drive you hom ne day of your proce	al or injectable) and/or a	contraception. Colono after the procedure. unit. You will not be a or bus, but only if you a	scopy able to drive are
					ns. • or Lime only), (4) Dulco	Jay tablata (1) 220 gran	n hattle of
				ed, Blue or Purple G		nax tablets, (1) 238 gran	n bottle of
			-	-	atorauc.,		
	Days Before Colo			(date)	. Dan dala ana khasas sa sa sa		
	ive a light breaktas include:	t, then drink only ci	ear liquids the r	est of the day. Clear	liquids are those you ca	in "see through". Exam	pies of clear
iiquius		Water					
		Strained fruit juice	without the pu	ıln			
		Plain coffee and to	-	•			
		Clear broth or box	•				
		7 UP, Sprite and G					
		•	-	d Propel Water) (Av	oid red, blue and purple	e)	
	✓	Kool-Aid or other	fruit flavored dr	inks (Avoid red, blue	e and purple)		
	✓	Plain Jell-O withou	ut added fruit or	toppings (Avoid red	d, blue and purple)		
	✓	Popsicles (Avoid r	ed, blue and pui	rple)			
In the	Evening , drink one	bottle of Magnesiu	ım Citrate (10 oı	unces) and immedia	tely after that drink 2 to	3 eight-ounce glasses o	of clear
liquid.							
	<mark>ay Before Colon</mark> e		On	(date)			
All Day	Continue with cle	ear liquids only. Clea	r liquids are tho	se you can "see thro	ough". No LIQUIDS that	are red, blue or purple.	•
	0 p.m. Take 4 Dulo						
At 4:0	-	_		es of Gatorade. Shal	ke well until all of the M	iralax powder is dissolve	ed.
	_	the Gatorade if des					
At 5:0	-	ounce glass of the G rinking clear fluids ι		ralax mixture every	20-30 minutes until it is	all gone.	
Day o	f Colonoscopy	=	On	(date)			

unless otherwise instructed by your physician. No gum or candy. **Medications** may be taken with sips of water on your regular schedule until 4 hours before your arrival time. **Diabetic patients:** please check your blood sugar level before your procedure.

You may have small amounts of clear liquids until 4 hours prior to your arrival time, then nothing to eat or drink until after the procedure,