



Lifespan

Delivering health with care.

Center for Women's Gastroenterology

Providence, RI

(401) 793-7080

Lifespan Physician's Group, Inc.

East Providence, RI

(401) 606-4260

Thomas DeNucci, MD

Fadlallah Habr, MD

Zilla Hussain, MD

Sarah Hyder, MD

Colleen Kelly, MD

Amanda Pressman, MD

Shannon Simmons, MD

Laura Massa, NP

Erica Taylor, PA

Your procedure will take place at:

- Rhode Island Hospital, 593 Eddy Street, Providence, RI - Main Building, 9th Floor
- Medical Office Center (MOC) Building, 2 Dudley Street, 3rd Floor, Providence, RI
- Wayland Surgical Center, 17 Seekonk Street, Providence, RI
- The Miriam Hospital, 164 Summit Avenue, Providence, RI
- Blackstone Surgical Center, 1526 Atwood Avenue, Suite #300, Johnston, RI

Instructions for Colonoscopy

On the day of your procedure please bring photo ID and a list of ALL your medications and dosages.

Due to unexpected emergencies, **your procedure time cannot be guaranteed.** Please contact your insurance company prior to your procedure to understand your responsibility for any co-payment and/or deductible. Note: This is a hospital-based Endoscopy Unit.

Date: _____ Appointment Time: _____ Arrival Time: _____ With: Dr. _____

Important

- Do not stop Aspirin unless instructed by your primary care physician or cardiologist. If you take an oral contraceptive at night, you should take the pill in the morning with clear liquid and use a backup method of contraception. Colonoscopy preparation may interfere with absorption of the pill. Resume your birth control pill as usual after the procedure.
- A responsible adult must be available to come into the Endoscopy suite to sign you out of the unit. You will not be able to drive on the day of your procedure. You must have someone drive you home. You may take a taxi or bus, but only if you are accompanied by a responsible adult.
- Please do not wear perfumes or body lotions on the day of your procedure.

Important: Notify your physician if you are taking medications for **diabetes (oral or injectable)** and/or **anticoagulants ("blood thinners")** such as **Coumadin, Warfarin, Heparin, Plavix**, etc., and ask for further instructions.

Purchase at the Pharmacy (2) Dulcolax (bisacodyl) tablets; and (2) 10 ounce bottles of Citrate of Magnesia (**lemon or lime flavor only**). Refrigerate the bottles of Citrate of Magnesia bottles if desired (this may help improve the taste).

Two Days Before Colonoscopy Avoid all popcorn, nuts, and seeds such as quinoa, sesame seeds, and whole flax seeds.

One Day Before Colonoscopy

You may have **liquids only throughout the day.**

NOTE: CLEAR LIQUIDS ONLY. AVOID liquids that are red, blue or purple.

Examples include: Water, Strained fruit juice without the pulp, Coffee, Tea, Clear broth or bouillon, 7 UP, Sprite and Ginger Ale, Sports drink (such as Gatorade and Propel Water) (Avoid red, blue and purple), Kool-Aid or other fruit flavored drinks (Avoid red, blue and purple), Plain Jell-O without added fruit or toppings (Avoid red, blue and purple), Popsicles (Avoid red, blue and purple)

Lunch: You may have milk and yogurt (even frozen), but it must be smooth in texture: no fruit, seeds, nuts, or solid pieces.

At 4:00 pm: Drink an 8 ounce glass of water or clear liquid.

At 5:00 pm: Drink (1) 10 ounce bottle of Citrate of Magnesia; and an 8 ounce glass of water or clear liquid.

At 7:00 pm: Drink an 8 ounce glass of water or clear liquid.

At 8:00 pm: Take (2) Dulcolax (bisacodyl) tablets with an 8 ounce glass of water or clear liquid.

At 9:00 pm: Drink an 8 ounce glass of water or clear liquid.

Day of Colonoscopy

At _____ am: Drink (1) 10 ounce bottle of Citrate of Magnesia.

You may have clear liquids up until 4 hours before your arrival time, then nothing by mouth until after the procedure unless otherwise instructed by your physician. **No gum or candy.** Medications may be taken with sips of water on your regular schedule until 4 hours before your arrival time. **Diabetic patients: please check blood sugar level before your procedure.**

The end result of your cleansing needs to be liquid stool, most often yellow in color. There may be small flecks of sediment in it.

Please call the office if you have any questions.